

PARENT RESOURCE GUIDE

Autism and Self-Stimulatory Behaviors

Understanding, purpose, impact, and treatment.

Dr. John Carosso, Psy.D.



The information in this guide is drawn from the clinical expertise of Dr. John Carosso, Psy.D. and Dr. Robert A. Lowenstein, M.D., current research findings, and authoritative sources including the DSM-5 and the National Institute of Mental Health. It is intended for informational purposes and does not replace a professional evaluation.

INTRODUCTION

Self-stimulatory behaviors, often referred to as “stimming,” are repetitive movements, sounds, or actions commonly observed in individuals with autism spectrum disorder (ASD). These behaviors are a hallmark feature of autism but are not exclusive to it – neurotypical individuals also engage in similar actions (such as tapping feet or twirling hair). For those with autism, however, stimming often serves unique and essential purposes.

WHAT ARE SELF-STIMULATORY BEHAVIORS?

Defining Stimming

Stimming includes a wide range of repetitive actions or movements that serve to regulate sensory input or emotional states. These behaviors are part of the diagnostic criteria for ASD under “restricted, repetitive patterns of behavior, interests, or activities” as outlined in the DSM-5.

Common examples of stimming:

- Physical movements: Flapping hands, rocking back and forth, spinning, or pacing.
- Verbal stims: Repeating words, phrases, or sounds.
- Tactile behaviors: Touching certain textures or tapping objects.
- Visual stimulation: Staring at lights, spinning objects, or watching repetitive movements.
- Auditory stimulation: Making clicking sounds, humming, or repeating music.

Why Do People Stim?

1. Self-regulation: Managing sensory input to reduce over- or under-stimulation.
2. Emotional expression: Coping with emotions such as excitement, anxiety, frustration, or boredom.
3. Sensory seeking: Gaining sensory feedback that feels pleasurable or calming.
4. Focus and concentration: Some individuals use stimming to enhance attention during tasks.

ARE SELF-STIMULATORY BEHAVIORS NORMAL?

Stimming is not exclusive to autism. Neurotypical individuals also engage in similar behaviors: tapping or fidgeting, hair twirling, chewing or biting, rhythmic movements, repetitive doodling, and verbal repetition. The key difference lies in the degree, frequency, and visibility of these behaviors, which tend to be less prominent in neurotypical individuals.



PURPOSE AND BENEFITS OF STIMMING

Sensory Processing and Regulation

Many individuals with autism experience sensory processing differences — hypersensitivity or hyposensitivity to sensory stimuli. Rocking back and forth might help calm an overstimulated individual, while flapping hands might provide needed stimulation in a dull environment.

Emotional Expression and Stress Relief

Stimming can act as an outlet for emotions, functioning as a self-soothing mechanism during periods of stress or excitement. A child may bounce or spin when excited, or hum softly to calm down in a loud and overwhelming space.

Communication Tool

In some cases, stimming serves as a nonverbal form of communication — signaling an individual's internal state and offering insight into how they feel or what they need.

Creativity and Focus

Certain stims can enhance focus and aid in creative problem-solving. Repetitive drawing or arranging objects can be calming and help the individual stay engaged in a task.

IMPACT OF STIMMING

Positive Impacts	Potential Challenges
1. Self-regulation — helping individuals manage emotions and sensory input. 2. Stress relief — offering a healthy outlet for overwhelming emotions. 3. Improved focus — assisting concentration during challenging tasks.	1. Safety concerns — behaviors like head-banging, biting, or hitting can lead to self-injury. 2. Social stigma — visible stimming may result in misunderstanding or bullying. 3. Interference with daily life — excessive stimming might disrupt schoolwork or social engagement.

Addressing these challenges requires a balanced approach that respects the individual's needs while ensuring their safety and well-being.

TREATMENT AND MANAGEMENT

When Is Treatment Needed?

Stimming should only be addressed if it poses a risk to the individual, significantly interferes with daily functioning or quality of life, or disrupts the individual's ability to learn or participate in desired activities.

The goal of treatment is not to eliminate stimming, but to support the individual in using it safely and adaptively.

Approaches to Treatment

- Behavioral Interventions (ABA): Identify the function of stimming and teach alternative behaviors to fulfill the same need.
- Sensory Integration Therapy: Weighted blankets or noise-canceling headphones may reduce the need for stimming in stressful environments.
- Occupational Therapy: Build self-regulation skills and find safe, socially appropriate ways to stim.
- Environmental Modifications: Dimming bright lights, creating quiet spaces, or ensuring predictable routines.
- Medication: In rare cases, to address underlying conditions like severe anxiety or ADHD. Generally a last resort.
- Parent and Caregiver Education: Fostering understanding and patience to create a supportive environment.

ABA AND SELF-STIMULATORY BEHAVIORS

ABA does not aim to eliminate stimming indiscriminately but to address behaviors that are harmful, stigmatizing, or disruptive. ABA practitioners begin with a Functional Behavior Assessment (FBA) to determine why the individual engages in stimming, then use reinforcement strategies including DRA, DRI, and DRO to teach replacement behaviors.

Example

A 7-year-old boy rocks during group activities. The FBA determines this helps him manage anxiety in noisy environments. A therapist introduces a weighted vest and deep-breathing exercises as alternatives. Data shows a significant reduction in rocking, and the boy now uses the weighted vest as a calming tool.

Ethical Considerations

ABA practitioners must balance addressing behaviors with respecting the individual's autonomy. Suppressing stimming without appropriate alternatives can increase anxiety and frustration. With growing advocacy for neurodiversity, ABA programs are increasingly adopting approaches that celebrate individuality and support safe, adaptive stimming.

SUPPORTING AUTISTIC INDIVIDUALS

Rather than discouraging stimming altogether, caregivers and educators can provide safe alternatives for harmful behaviors, offer sensory-friendly tools like chewable necklaces or fidget spinners, and normalize stimming in inclusive settings to reduce stigma.

Conclusion

Self-stimulatory behaviors are a fundamental aspect of life for many individuals with autism, serving purposes such as sensory regulation, emotional expression, and focus. Embracing a compassionate, individualized approach that balances safety, autonomy, and acceptance is essential. By fostering understanding and respecting the role stimming plays, we can support autistic individuals in leading fulfilling lives.

Ready to take the next step?

Doctors Carosso and Lowenstein and our experienced team are here to help. Contact us to schedule an evaluation or to ask any questions.
(724) 733-5757 | hello@acpitt.com | acpitt.com